Cosmology from Home 2021 5 July 2021 – 16 July 2021

Probing Interacting Dark Energy and Scattering of Baryons with Dark Matter in Light of EDGES 21cm Signal

Upala Mukhopadhyay

Astroparticle Physics and Cosmology Division

Saha Institute of Nuclear Physics, Kolkata, India.

Collaborators: Debasish Majumdar, Kanan K. Datta; Phys.Rev.D 103 (2021) 6, 063510

Plan of Talk


Objective or Motivation


Formalism


Results

Conclusion and take home messages

21cm Line and Cosmic Dawn







we observe 21cm line with CMB as background light


$${
m if} \; T_s > T_{
m cmb} \;\;\;\;\; {
m emission}$$
 ${
m if} \; T_s < T_{
m cmb} \;\;\;\;\; {
m absorption}$

Because it is a line, we can observe the universe at each redshift

redshift	λ	ν
0	21cm	1420MHz
9	2.1 m	142MHz
19	4.2 m	71MHz

Spin Temperature


- The excitation temperature of 21 cm line is known as the spin temperature, $n_1/n_0 = g_1/g_0 \exp(-T_*/Ts)$
- Depends on i) CMB ii) gas temperature iii) Lyα photons



Problems in EDGES Results

Large absorption feature than expected

Smaller spin temperature,
Larger background
temperature,
Larger optical depth




Doi:10.1038/nature25792

Problems in EDGES Results

Large absorption feature
than expected

•Dark Matter – Baryon Interaction
•Interacting Dark Energy scenario

Constraints on model parameters



Doi:10.1038/nature25792

Effects of the Interactions Between DM and Baryon Fluid

- There is a temperature difference as well as velocity difference between the DM and baryon fluid.
- The interactions between two fluids of different temperature will heat up the colder fluid and cool down the warmer one. Also for the different velocities, a heating term can arise.

$$\frac{dQ_b}{dt} = \frac{2m_b \rho_{\chi} \sigma_0 e^{-r^2/2} (T_{\chi} - T_b)}{(m_{\chi} + m_b)^2 \sqrt{2\pi} u_{\text{th}}^3} + \frac{\rho_{\chi}}{\rho_m} \frac{m_{\chi} m_b}{m_{\chi} + m_b} V_{\chi b} D(V_{\chi b})$$

The dragging term to damp the relative velocity

$$D(V_{\chi b}) \equiv -\frac{dV_{\chi b}}{dt} = \frac{\rho_m \sigma_0}{m_b + m_\chi} \frac{1}{V_{\chi b}^2} F(r)$$

$$F(r) \equiv \operatorname{erf}(\frac{r}{\sqrt{2}}) - \sqrt{\frac{2}{\pi}} e^{-r^2/2} r$$

$$r \propto V_{\chi b}$$

J. B. Munoz, et.al, Phys. Rev. D 92, 083528 (2015).

Effects of Interactions Between DM and DE on 21cm Absorption Line

 Considering the interactions between DM and DE, the continuity equations are given as

$$(1+z)H(z)\frac{d\rho_{\chi}}{dz} - 3H(z)\rho_{\chi} = -Q ,$$

$$(1+z)H(z)\frac{d\rho_{de}}{dz} - 3H(z)(1+\omega)\rho_{de} = Q$$

Modification of the evolution of the Universe due to this interaction

$$H(z) \neq H_0 \sqrt{\Omega_{m0}(1+z)^3 + \Omega_{de0}(1+z)^{3(1+\omega)}}$$

It will modify the optical depth

$$\tau = \frac{3}{32\pi} \frac{T_*}{T_s} n_{\rm HI} \lambda_{21}^3 \frac{A_{10}}{H(z)}$$

DM-DE interaction (contd.)

Three phenomenological models of this interaction are

$$M-I$$
 Q=3 $\lambda H(z)\rho_{\rm de}$,
 $M-II$ Q=3 $\lambda H(z)\rho_{\chi}$,
 $M-III$ Q=3 $\lambda H(z)(\rho_{\rm de}+\rho_{\chi})$

Experimental constraints on these models

Model	ω	λ	H_0
$3\lambda H \rho_{ m de}$	$-0.9191^{+0.0222}_{-0.0839}$	$-0.1107^{+0.085}_{-0.0506}$	$68.18^{+1.43}_{-1.44}$
$3\lambda H \rho_{ m de}$	$-1.088^{+0.0651}_{-0.0448}$	$0.05219^{+0.0349}_{-0.0355}$	$68.35^{+1.47}_{-1.46}$
$3\lambda H \rho_{\chi}$	$-1.1041^{+0.0467}_{-0.0292}$	$0.0007127^{+0.000256}_{-0.000633}$	$68.91^{+0.875}_{-0.997}$
$3\lambda H(ho_{ m de}+ ho_\chi)$	$-1.105^{+0.0468}_{-0.0288}$	$0.000735^{+0.000254}_{-0.000679}$	$68.88^{+0.854}_{-0.97}$

C. Li, et.al, Phys. Lett. B801 (2020) 135141.

 We will investigate that whether a IDE model, which is well in agreement with the constraints given from other experiments, could also be consistent in explaining the EDGES results.

Equations to Solve

Temperature evolution of DM

$$\frac{dT_{\chi}}{dz} = \frac{2T_{\chi}}{1+z} - \frac{2\dot{Q}_{\chi}}{3H(1+z)} - \frac{1}{n_{\chi}} \frac{2Q}{3H(1+z)}$$

Temperature evolution of baryon

$$\frac{dT_b}{dz} = \frac{2T_b}{1+z} + \frac{\Gamma_c}{H(1+z)}(T_b - T_\gamma) - \frac{2\dot{Q}_b}{3H(1+z)}$$

Evolution of free electron fraction

$$\frac{dx_e}{dz} = \frac{C_P}{H(1+z)} \left(n_H A_B x_e^2 - 4(1-x_e) B_B e^{\frac{-3E_0}{4T_\gamma}} \right)$$


Variation of relative velocity

$$\frac{dV_{\chi b}}{dz} = \frac{V_{\chi b}}{1+z} + \frac{D(V_{\chi b})}{H(1+z)}$$

Brightness temperature of 21 cm line


$$T_{21} = \frac{T_s - T_{\gamma}}{1+z} (1 - \exp^{-\tau}) \approx \frac{T_s - T_{\gamma}}{1+z} \tau$$
 $T_b = T_s$

Hubble Parameter




21 cm Brightness Temperature for Model-I for Different Cases

• EDGES limit $-0.1 \text{K} \ge \Delta T_{21} \ge -0.8 \text{K}$, where $\Delta T_{21} = (T_{21} - T_{21}^0)$.





21 cm Brightness Temperature for Model-III for Different Cases

- EDGES observations are satisfied for interaction strength greater or equal to 0.04.
- It is shown from other experiments that λ should not be greater than 0.000989.
- Same conclusions can be obtained for Model-II.




Allowed Parameter Space





Allowed Parameter Space

$$-300 \text{mK} \ge T_{21} \ge -1000 \text{mK}$$



Comments, We Can make

 The EDGES experiment has observed an excess trough in the brightness temperature of the 21cm absorption line

Baryons-DM interaction

DM-DE interaction

- Larger DM-baryon interaction cross section, larger DM-DE interaction parameter and smaller DM mass are more favourable to achieve the excess absorption feature.
- When Model-I is considered for DM-DE interaction, EDGES results and other experiment results are well respected but it is not so for other two IDE models.
- DM-DE interaction raises the possibility of probing larger mass ranges of DM that could have influenced the cooling effects.

THANK YOU